

# W/Z + jets in ATLAS

**Pierre-Hugues Beauchemin**

**University of Oxford**



**On behalf of the ATLAS Collaboration**



Standard Model Benchmark Workshop at the Tevatron and the LHC



Fermilab, November 19<sup>th</sup>-20<sup>th</sup>, 2010

# Scope of the presentation

- Level of precision needed on  $W/Z$ +jets measurements for sensitivity to improvement in theoretical understanding of these processes
- State where  $W/Z$ +jets ATLAS measurements stand and show data to MC comparison of various observables
- Give a quick overview of how we tackle down some of the important systematics affecting  $W/Z$ +jets analyses
- Bring to your attention some issues we are facing in order to start useful discussions
  - Bring feed-back to the collaboration
  - Hopefully reach some consensus among the wide HEP community

# W/Z+jets measurements


- ATLAS is in commissioning period:
  - Jet &  $E_T^{\text{miss}}$  resolution and calibration
  - Leptons energy scale and resolution
  - Trigger, Pile-up and Luminosity
- Crucial to study W/Z+jets events to understand our detector and tools
  - First priority with 2010 data




Perturbative QCD

Fragmentation  
Hadronization  
Underlying events

The key to reach a better understanding of the Standard Model is:  
**to keep systematic uncertainties low**



# Status of W/Z+jets analyses



- We studied W+jets and Z+jets with 1 pb<sup>-1</sup>
  - Absolute cross sections
  - Relative to inclusive cross sections
  - All possible ratios

Uncertainty on W+jets already dominated by systematic uncertainty

- MC-based or simplify correction factors
- Mostly conservative estimate of systematic uncertainties
- Paper under internal review

- Z+jets uncertainty :  $\Delta_{\text{stats}}$    $\Delta_{\text{syst}}$ 
  - Public note under internal review

- Provide Comparison to LO and NLO (MCFM) calculations
- Update with 2010 full 45 pb<sup>-1</sup> recorded dataset
  - Use more data-driven corrections

# **FIRST LOOK AT W/Z+JETS ATLAS DATA**

# Data vs MC

- Use 35 to 42 pb<sup>-1</sup> ( $\pm 11\%$ ) of data in the following distributions
  - Collected from 6 GeV to 15 GeV thresholds electron and muon triggers
- MC used in the following distributions
  - ALPGEN+HERWIG+JIMMY with CTEQ6L1 PDF for W/Z+jets events
  - Pythia dijet events ( $P_T > 15$  GeV) with MRST2007LO\* PDF for QCD
  - Use POWHEG with CTEQ6L1 PDF for ttbar
  - Added  $\langle N \rangle = 2$  pile-up events, reweighted to primary vertices observed in data
  - ATLAS MC09 tune are used
- MC events normalized to observed data candidates before jet selections
  - Relative normalization of MC samples to NLO cross sections except QCD


# Selections

- Lepton kinematic:
  - Electron  $E_T^{\text{clus}} > 20$  GeV and muon combined track  $P_T > 20$  GeV with  $P_T^{\text{MS}} > 10$  GeV
- Eta coverage:
  - $|\eta_{\text{ele}}| < 2.47$  excluding barrel to end-cap transition region ( $1.37 < |\eta_{\text{ele}}| < 1.56$ ),  
 $|\eta_{\text{muo}}| < 2.4$
- Lepton quality requirements:
  - Tight requirements on electron cluster shape, track quality and matching
  - Muon cone 4 track isolation  $\Sigma P_T^{\text{ID}}/P_T < 0.2$  and  $|P_T^{\text{ID}} - P_T^{\text{MS}}| < 15$  GeV
- Jet selections:
  - AntiKt4 jets with  $P_T > 20$  GeV and  $|\eta| < 2.8$
- ETmiss selections
  - $E_T^{\text{miss}} > 25$  GeV, computed from calibrated topoclusters and out-of-cluster energy
- ATLAS standard clean-up cuts are applied


# Jet Multiplicity

- QCD and top backgrounds significantly increase with the number of jets
  - Bigger effect at higher centre of mass energy
- ⇒ Need to estimate these backgrounds precisely from data
- Other electroweak backgrounds can be estimated using MC ratios

Electron channel



Muon channel




Remember:  
jet threshold  
is  $E_T > 20 \text{ GeV}$




# Lepton Transverse Momentum

- MC describes well the detector effects on lepton reconstruction and resolution
- Can use MC to estimate electron and muon resolution effects on acceptance
  - For better precision, measure resolution in data before correction.

Electron channel







Muon channel




# Jet Transverse Momentum

- Simulation are used to unfold detector effects in W/Z+jets measurements
  - ATLAS data well modelled by simulation:
    - Jet transverse momentum in W+jets events
    - reconstruction efficiency in QCD dijet events
    - jet energy resolution in QCD dijet events
- Will be measured on Z+jets events



# Missing $E_T$


- Reasonably good agreement between data and MC in the bulk of the signal region
  - some care must be made with missing  $E_T$  model
    - pile-up effects
- Use data to estimate:
  - Missing  $E_T$  selection acceptance correction
  - QCD background prediction



Electron channel

Low pile-up


Higher pile-up




# W/Z Transverse Momentum

- Good understanding of vector boson transverse momentum reconstructed from leptons over wide range of kinematics
  - Will be used to calibrate jets in W/Z+jets events
  - Useful to tune soft QCD effects in MC

**W+≥1-jet  
Electron channel**





**Z+≥0-jet  
Muon channel**



# **TOWARD PRECISE W/Z+JETS MEASUREMENTS**


# Systematic: QCD background

- QCD bkg predictions proceed from the opposition of two big numbers
  - Dijet cross section
  - Fake rejection
  - ⇒ Hard to estimate from MC
- Template method ( $W \rightarrow ev$ ):
  - QCD  $E_T^{\text{miss}}$  from reversed electron selections,  $W E_T^{\text{miss}}$  from MC
  - Fit both templates to data
  - Integrate normalized QCD in signal region
  - ▶  $f_{\text{QCD}}(W+\geq 1\text{-jet}) \sim 12\%$ ,  $\Delta f_{\text{QCD}}/f_{\text{QCD}} \sim 30\%$
- $Z \rightarrow ee$  QCD background estimate:
  - Direct fit to invariant mass
  - Number of same sign leptons under Z peak
  - ▶  $f_{\text{QCD}}(Z+\geq 1\text{-jet}) \sim 3\%$ ,  $\Delta f_{\text{QCD}}/f_{\text{QCD}} \sim 30\%$



# Lepton efficiency

- Lepton efficiencies don't depend on the recoiling jet activity
  - Can use precise estimates from inclusive data samples
- Efficiencies are measured from MC in first 2010 analyses
  - Tag & Probe method and  $E_T^{\text{miss}}$  preselected events already show promising results
- Trigger fully efficient in 2010 data



Parameter	$\delta C_W / C_W (\%)$	$\delta C_Z / C_Z (\%)$
Trigger efficiency	<0.2	<0.2
Material effects, reconstruction and identification	5.6	8.8
Energy scale and resolution	3.3	1.9
$E_T^{\text{miss}}$ scale and resolution	2.0	-
Problematic regions in the calorimeter	1.4	2.7
Pile-up	0.5	0.2
Charge misidentification	0.5	0.5
FSR modelling	0.3	0.3
Theoretical uncertainty (PDFs)	0.3	0.3
Total uncertainty	7.0	9.4


$$C_W = \epsilon_{\text{event}}^W \cdot \alpha_{\text{reco}}^W \cdot \epsilon_{\text{lep}}^W \cdot \epsilon_{\text{trig}}^W$$

$$C_Z = \epsilon_{\text{event}}^Z \cdot \alpha_{\text{reco}}^Z \cdot (\epsilon_{\text{lep}}^Z)^2 \cdot [1 - (1 - \epsilon_{\text{trig}}^Z)^2]$$


Eliminated from data-driven estimate of efficiencies

# Systematics: $E_T^{\text{miss}}$

- Need to correct for detector effects on  $E_T^{\text{miss}}$  selection acceptance
  - Non-trivial systematic uncertainty
    - Jet energy scale and resolution
    - Pile-up
    - Material modelling
    - Non-cluster energy
- $\Rightarrow \Delta A_{\text{reco}} \sim 2\text{-}4\%$  from MC estimate



- $E_T^{\text{miss}}$  resolution can be measured in data




Data-driven estimate of correction factor to  $E_T^{\text{miss}}$  acceptance

- Select Z+jets events
- Measure  $\sigma(E_T^{\text{miss}})$  along  $n_{\text{perp}}$
- Measure  $\mu(E_T^{\text{miss}})$  along  $n_{\text{para}}$
- Apply gaussian smearing to true  $P_T^{\nu}$



# Systematics: detector unfolding

- Jet energy resolution and reconstruction well modelled in MC.
- To correct measurement up to hadron level, need to solve the reverse problem (unfolding)
  - involve other sources of uncertainties
  - Many different techniques on the market
  - More complicated problem
- Smaller than lepton efficiency correction
  - Not a dominant source of systematics in 2010 measurements
  - Used simple bin-by-bin corrections
- Need to adopt:
  - good unfolding method,
  - generate high MC statistics,
  - identify all sources of systematics






Unfolding workshop at CERN

on 20/01/2011

# Systematics: Jet Energy Scale

- ATLAS jets are calibrated from MC
  - Conservative uncertainty estimate
- Large impact of Jet Energy Scale uncertainty on cross sections
  - $\Delta\text{JES} = 5\% \rightarrow \Delta\sigma_{\text{tot}} \sim 7\%$
  - Remove statistical effects in  $\Delta\sigma$
  - Smaller effects on ratios
- Single particle measurement will constrain  $\Delta\text{JES}$  in 2011 analyses



# Systematics: summary and outlook

- Dominant systematic uncertainties on W/Z+1-jet  $\sim 1\text{pb}^{-1}$  analysis

Sources of systematics	$\Delta\sigma(\text{W+jets}) 1 \text{ pb}^{-1}$	$\Delta\sigma(\text{Z+jets}) 1 \text{ pb}^{-1}$	target for $45\text{pb}^{-1}$ analyses
Jet Energy Scale	10%	10%	6-7%
Lepton $A_{\text{det}}$ and $\varepsilon$	7%	10%	4-5%
Unfolding + jet $E_{\text{T}}$ resolution	3%	5%	3%
QCD background	6%	2%	1-3%
$E_{\text{T}}^{\text{miss}} A_{\text{det}}$	3%	-	1-2%
Luminosity	11%	11%	5%

- Total of  $\sim 14/15\%$  systematic uncertainty (excluding luminosity) compared to 3/11% statistical uncertainty
  - Already systematic dominated with  $\sim 1\text{pb}^{-1}$  of data
  - Factor of  $\sim 2$  reduction on systematic and luminosity uncertainties with full dataset
- ⇒ Sensitivity to NLO effects
- ⇒ Will soon start to get sensitivity beyond theoretical precision

Effects which were small and neglected must now be treated properly

⇒ Induce some discussions


# DISCUSSION

# Jet Energy Threshold

- Experimental criteria favour jet  $P_T$  thresholds above 30 GeV:
  - Calibration below 20 GeV affected by jet reconstruction threshold
  - Jet energy scale uncertainty and pile-up dramatically increase below 30 GeV
  - Jet reconstruction efficiency quickly decreases below 30 GeV
- Theoretical prediction less robust for lower jet  $E_T$ :
  - Low  $P_T$  jets are more sensitive to soft QCD effects
  - Lower  $P_T$  jet thresholds imply higher scale uncertainty for theoretical predictions

Q: Prefer to reach as low jet  $P_T$  thresholds as possible (input to theorists) or keep thresholds higher (more robust comparisons)?

Q: What would be desirable jet thresholds in W/Z+jets physics?



# Detector acceptance

- Lepton acceptance corrections depend on theoretical input (generators)
  - True level lepton  $P_T$  and  $\eta$  cuts,  $E_T^{\text{miss}}$  cut, mass selections, etc
- ⇒ Acceptance correction in measurements make it difficult for theorist to disentangle these effects to test potential improvements in their models
- Correcting for detector acceptance is needed to:
  - Combine or compare measurements made in muon and electron channels
  - Compare results from various experiments (detector independent results)

Q: Prefer to see publication with visible cross sections only or correct for a full acceptance cross sections?

Q: How isolation should be treated?

So far, in ATLAS, we are working out both numbers, but publication preferences still on analysis to analysis basis.

# QED Final State Radiation

- QED radiation, especially from FSR is:
  - different for electrons and muons
  - simulated with varying accuracy in different MC programs and kinematic regions
- Theory accuracy at the few-per-mille level for inclusive cross section,
  - ▶ this is *not true* in general for differential or exclusive distributions.

Final state electron	Pythia status code 1	Well defined physics final state	Input to unfolding is theory dependent
Electron at production vertex	Pythia status code 3	Compare electrons and muons	Measurement can't profit from new theory
"Dressed" electron	Final state + photon in cone around it	Physics final state close to vertex electron	Cone size is arbitrary


Q: what exactly should we measure and how should we confront it with theory? Correction before or after radiations? Both?

Q: How to assign reasonable systematic uncertainty on such effect?

Q: Should QED radiation be included in true jet clustering?

# Jet-Electron overlap

- Electrons are reconstructed as jets in the calorimeter
  - Affect the energy response and reconstruction efficiency of close-by jets
- No ideal way to experimentally deal with this
  - Remove jets using  $\Delta R$  matching
    - Small cone ( $\sim 0.2$ )
    - Large cone ( $\sim 0.5$ )
    - Small cone + event veto
  - Remove electron cluster from calorimeter and rerun jet algorithm
  - 4-vector subtraction




Q: Which is the best approach for comparison to theory and with other experiments?

Q: When a  $\Delta R$  approach is used, should the decay products of vector boson be included in true jet clustering?



# Correction to Parton level MC

- Unfolding brings jets from detector level to hadron level
- NLO fixed order calculation programs like MCFM don't include non-perturbative QCD effects for W/Z+parton(s) processes
  - Hadronization, underlying event
    - ⇒ No hadron level...
- K-factors obtained from NLO/LO MCFM prediction can't be applied to predictions involving parton shower



Q: Should we compute hadronization + underlying event correction factor from PYTHIA or HERWIG and apply them to MCFM predictions or it is preferable to leave predictions as they are?

# Conclusion

- W/Z+jets physics allow to:
  - Study detector performance and validate measurement tools
    - ⇒ Improve systematic uncertainty on calibrations, efficiencies, unfolding, etc
  - Better understanding of higher order QCD corrections
    - ↻ Require small systematic uncertainty on the measurements

Provide crucial understanding of major background to many new physics searches

- ATLAS performed 1 pb<sup>-1</sup> W/Z+jets measurements
  - Good data to MC agreement
  - Set the ground for more precise future measurements
- Small effects will become important as the precision increases
  - ▶ Need already discussions with theorist and other experiments on how to provide the best handle on these effects
    - Eg: QED FSR, jet energy threshold, jet-electron overlap removal, etc
- Using the 2010 full dataset and the yet to come 2011 data:
  - improve jet,  $E_T^{\text{miss}}$  and lepton performances using Z+jets events
  - Start to study Heavy flavour

# Some references

- First Z+jets MC study:
  - CSC book: [arXiv:0901.0512](#) [hep-ex]
- Paper on the W/Z inclusive measurement
  - CERN-PH-EP-2010-037, [arXiv:1010.2130](#) [hep-ex]
- Jet energy resolution and reconstruction efficiency studies
  - [ATLAS-CONF-2010-054](#)
- Jet energy scale uncertainty estimate
  - [ATLAS-CONF-2010-056](#)
- ETmiss performance studies
  - [ATLAS-CONF-2010-057](#)