

Rare and Radiative Kaon Decays from the NA48/2 Experiment

Vladimir Kekelidze
(JINR, Dubna)

on behalf of the **NA48/2** Collaboration:
Cambridge, CERN, Chicago, Dubna, Edinburgh,
Ferrara, Firenze, Mainz, Northwestern, Perugia, Pisa, Saclay,
Siegen, Torino, Vienna




Content

- **The NA48 detector**
- **$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$ (Ke4)**
- **$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$**
- **$K^\pm \rightarrow \pi^\pm \gamma \gamma$**
- **Conclusions**


The NA48 detector

Main detector components:

- **Magnetic spectrometer (4 DCHs):**
4 views/DCH:
 $\Delta p/p = 1.0\% + 0.044\% * p [GeV/c]$.
- **Hodoscope**
fast trigger (150ps).
- **Liquid Krypton EM calorimeter (LKr)**
high granularity, quasi-homogeneous;
 $\sigma_E/E = 3.2\%/E^{1/2} + 9\%/E + 0.42\% [GeV]$;
 $\sigma_x = \sigma_y = 0.42/E^{1/2} + 0.6mm (1.5mm @ 10GeV)$
- **Hadron calorimeter**
- **Muon veto counters**
- **Photon vetoes**



2-3M K/spill ($\pi/K \sim 10$),
 π decay products stay in pipe;
 flux ratio: $K^+/K^- \approx 1.8$



K^\pm beams

$$\mathbf{K}^{\pm} \longrightarrow \pi^{+} \pi^{-} e^{\pm} \nu$$

1.13 M decays

The measurements of:


- Form-Factors
- pion-pion scattering lengths: \mathbf{a}_0 and \mathbf{a}_2

Ke4 decays : formalism

Five kinematic variables (Ca.Ma. variables)

(Cabibbo-Maksymowicz 1965)

$S_\pi (M^2_{\pi\pi})$, $S_e (M^2_{e\nu})$,
 $\cos\theta_\pi$, $\cos\theta_e$ and ϕ .



Partial Wave expansion of the amplitude

into s and p waves (Pais-Treiman 1968)

+ Watson theorem (T-invariance) for δ_l^I

$$\delta_0^0 \equiv \delta_s \text{ and } \delta_1^1 \equiv \delta_p$$

F, G = 2 Axial Form Factors

$$F = F_s e^{i\delta_s} + F_p e^{i\delta_p} \cos\theta_\pi$$

$$G = G_p e^{i\delta_g}$$

H = 1 Vector Form Factor

$$H = H_p e^{i\delta_h}$$

F, G, H are complex and dimensionless

Map the distributions of the Ca.Ma. variables in the **five-dimensional space** with 4 Form factors and one phase shift, assuming identical phases for the p-wave Form Factors F_p, G_p, H_p :

The fit parameters are :

$$F_s, F_p, G_p, H_p \text{ and } \delta = \delta_s - \delta_p$$

(F_s, F_p, G_p, H_p are real)

Ke4 selection & background

Signal ($\pi^+ \pi^- e^\pm \nu$) topology:

- 3 charged tracks, good vertex
- 2 opposite sign pions
- 1 electron ($E_{LKr} / P \sim 1$)

Main background sources:


a) $K^+ \rightarrow [\pi^- \rightarrow e^- \nu] \pi^+ \pi^+$

b) $K^+ \rightarrow [\pi^0 \rightarrow e^+ e^- (\gamma)] (\pi^0) \pi^+$

Background seen in «wrong sign» (WS) component (same sign pions)

a) **1**(WS) / **2**(RS)

b) **1**(WS) / **1**(RS)



Total background is at the level of **2 x 0.3 %**
(estimated from WS events & checked by MC)

1.13 M of fully reconstructed **Ke4** (2003+2004 data)

Ke4 formfactor fit results

Series expansion with:

- $q^2 = S\pi/4m_\pi^2 - 1$
- $Se/4m_\pi^2$

$$F_s^2 = f_s^2(1 + f'_s/f_s q^2 + f''_s/f_s q^4 + f'_e/f_s Se/4m_\pi^2)^2$$

$$G_p/f_s = g'_p/f_s + g''_p/f_s q^2$$

Preliminary
(2003+2004)

	value	stat	syst
f'_s / f_s	0.152	± 0.007	± 0.005
f''_s / f_s	-0.073	± 0.007	± 0.006
f'_e / f_s	0.068	± 0.006	± 0.007
f'_p / f_s	-0.048	± 0.003	± 0.004
constant			
g_p / f_s	0.868	± 0.010	± 0.010
g'_p / f_s	0.089	± 0.017	± 0.013
h_p / f_s	-0.398	± 0.015	± 0.008
constant			

From phase shifts to pion-pion scattering lengths: a_0 and a_2

- Roy equations (unitarity, analyticity & crossing symmetry)
numerical solutions compute phases from data
using a_0 and a_2 , in the *isospin symmetry limit*

- Parameterization of Roy equations numerical solution:


$$\tan(\delta_l^I(s)) = \sigma_\pi q^2 (A_l^I + B_l^I q^2 + C_l^I q^4 + D_l^I q^6) (4m_\pi^2 - s_l^I) / (s - s_l^I)$$

δ_l^I -phases; $A_l^I, B_l^I, C_l^I, D_l^I, s_l^I$ – coefficients,
each is expressed as a 3-rd degree polynomial expansion of a_0 & a_2
ACGL PR 353(2001)207; DFGS EPJ C24(2002)469.

- Isospin corrections are evaluated as a correction to
phase differences (*CGR EPJ C59(2009)777*)
- Radiative effects are included in simulation
— *Gamov factor ($\pi^+\pi$ Coulomb attraction) in matrix element
& PHOTOS generator for real photons emission*

Ke4 phase shift measurement

isospin correction (10-15 mrad) to all published points — downward shift errors — combined stat. and syst.; fit to NA48/2 data alone



Systematic errors:

-bin to bin correlated:


- background level;
- isospin corrections

- bin to bin uncorrelated:

- fitting procedure;
- trigger efficiency;
- acceptance control;
- background shape;
- electron identification;
- radiative correction

Ke4 phase shift measurement

isospin correction (10-15 mrad) to all published points — downward shift errors — combined stat. and syst.; fit to NA48/2 data alone



Systematic errors:

-bin to bin correlated:

- background level;
- isospin corrections

- bin to bin uncorrelated:

- fitting procedure;
- trigger efficiency;
- acceptance control;
- background shape;
- electron identification;
- radiative correction

Scattering lengths **preliminary** result

- 2 parameter fit (correlation 0.967)

$$a_0 = 0.2220 \pm 0.0128(\text{stat}) \pm 0.0050(\text{syst}) \pm 0.0037(\text{th}^*)$$

$$a_2 = -0.0432 \pm 0.0086(\text{stat}) \pm 0.0034(\text{syst}) \pm 0.0028(\text{th}^*)$$

- 1 parameter fit using ChPT link between a_0 and a_2

$$a_0 = 0.2206 \pm 0.0049(\text{stat}) \pm 0.0018(\text{syst}) \pm 0.0064(\text{th}^*)$$

* Theoretical errors evaluated from isospin corrections & inputs to Roy equations numerical solution


CGR EPJ C59 (2009) 777

Perfect agreement
with theory prediction:


$$a_0 = 0.220 \pm 0.005$$

$$a_2 = -0.0444 \pm 0.0010$$

CGL NP B603(2001), PRL 86 (2001)



Comparison with other measurements of a_0



$$\mathbf{K}^{\pm} \longrightarrow \pi^{\pm} \pi^0 \gamma$$



The measurements of:

- DE & INT terms in decay 600k decays
- CPV test 1M decays

$$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$$

3 terms: IB, DE & INT

contributing to the decay
could be disentangled in
W spectra
(integrated over $T^*\pi$)



$$W^2 = \frac{(P_\pi^* \cdot P_\gamma^*)(P_K^* \cdot P_\gamma^*)}{(m_K m_\pi)^2}$$

$$\frac{\partial \Gamma^\pm}{\partial W} = \underbrace{\frac{\partial \Gamma_{IB}^\pm}{\partial W}}_{\text{IB}} \left[1 + \underbrace{2 \cos(\pm \phi + \delta_1^1 - \delta_0^2) m_\pi^2 m_K^2 |X_E|}_{\text{INT}} W^2 + \underbrace{m_\pi^4 m_K^4 (|X_E|^2 + |X_M|^2)}_{\text{DE}} W^4 \right]$$

IB can be calculated as $\text{BR}(\pi^+\pi^0) + \text{QED corrections}$

DE $\mathcal{O}(p^4)$ contribution has 2 terms:

- X_M : magnetic part 2 contributions by chiral anomaly
- Reducible: calculated using WZW functional ($X_M \sim 270 \text{ GeV}^4$)
- Direct: cannot be calculated in model independent way
- X_E : no prediction in ChPT depends on unknown constants


INT arises from interference of **IB** and X_E **DE**

- no model independent prediction available in ChPT

$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$: event selection



- At least 1 track + 3 clusters in LKr;
- Acceptance and BG-rejection cuts (M_K , COG etc...);
- $T_\pi^* < 80$ MeV.

BG channel	BG mechanism
$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$	missing or overlapping γ
$K^\pm \rightarrow \pi^\pm \pi^0$	γ from accidental or hadronic shower
$K^\pm \rightarrow \pi^0 e^\pm \nu$	e mistagged as π , accidental γ
$K^\pm \rightarrow \pi^0 \mu^\pm \nu$	μ mistagged as π , accidental γ



Residual BG < 1%

Extended maximum likelihood fit



Likelihood fit full data sample

$$\text{Frac}(\text{DE}) = (3.32 \pm 0.15_{\text{stat}} \pm 0.14_{\text{syst}}) \cdot 10^{-2}$$

$$\text{Frac}(\text{INT}) = (-2.35 \pm 0.35_{\text{stat}} \pm 0.39_{\text{syst}}) \cdot 10^{-2}$$

First evidence of nonzero **INT** contribution

(systematics dominated by trigger efficiency)

PDG ($55 < T_{\pi^*} < 90$ MeV):

$$\text{Br} = (2.75 \pm 0.15) \cdot 10^{-4}, \quad [\text{DE} / \text{Br} = (1.56 \pm 0.27) \cdot 10^{-2}]; \quad \text{INT} - \text{no data}$$

$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ CPV tests

1) Difference in K^+ and K^- decay rates:

$$A_N = (N^+ - RN^-)/(N^+ + RN^-),$$

where $R(K^+/K^-)$ ratio = 1.7998 ± 0.0004 from $K3\pi$

Result :

$$A_N = -0.0 \pm 0.001_{\text{stat}} \pm 0.0006_{\text{syst}}$$


systematics from trigger and kaon momentum distributions

2) Difference in W spectra shapes:

$$\begin{aligned} dA_W/dW &= (\Gamma^+ - \Gamma^-)/(\Gamma^+ + \Gamma^-) = \\ &= eW^2/(1+aW^2+bW^4), \\ &e \text{ is a free parameter} \end{aligned}$$

Result (integrating):

$$(I_{\text{INT}}/I_{\text{IB}})A_W = -0.0006 \pm 0.0010_{\text{stat}}$$



$$\mathbf{K}^{\pm} \rightarrow \pi^{\pm} \gamma \gamma$$

1164 decays

The BR measurement

$K^\pm \rightarrow \pi^\pm \gamma\gamma$: Theory

$$\frac{\partial^2 \Gamma}{\partial y \partial z} = \frac{m_{K^+}}{(8\pi)^3} \left[z^2 (|A + B|^2 + |C|^2) + \left(y^2 - \frac{1}{4} \lambda(1, r_\pi^2, z) \right)^2 (|B|^2 + |D|^2) \right]$$

$$y = \frac{P \cdot (q_1 - q_2)}{M_K^2} \quad z = \frac{(q_1 + q_2)^2}{M_K^2} = \frac{M_{\gamma\gamma}^2}{M_K^2}$$

P =pion 4-momentum
 $q_{1,2}$ =photon 4-momenta

A, B, C, D depend on Z and on ChPT \hat{c} parameter




$\hat{c} = 0.90 \pm 0.45$ from $K^\pm \rightarrow \pi^\pm e^+ e^- \gamma$

ChPT contributions to the amplitude:

$O(p^4)$: $A(z, \hat{c})$ responsible for cusp at $m_{\gamma\gamma} = 2m_\pi$
 $C \neq 0$ for WZW anomaly (poles & tadpoles), $B = D = 0$

[D'Ambrosio, Portolés, PLB386 (1996) 403]

$O(p^6)$: unitarity corrections can increase BR by 30-40% (depending on \hat{c} value)



$K^\pm \rightarrow \pi^\pm \gamma\gamma$ Br measurement

Event selection

- At least 1 track and 2 LKr clusters
- COG, CDA vertex cuts
- $M_{\gamma\gamma}$ in kinematical limits

1164 events selected

20% of full data sample

- **40 times** more than the world sample
- **3% Bkg** mainly from $\pi^\pm \pi^0 \gamma$
[$Br = (2.75 \pm 0.15) \cdot 10^{-4}$]
- Normalization — $K2\pi$ (~6 M)
- Main systematics : **acceptance**



Preliminary:

$$Br = (1.07 \pm 0.04_{\text{stat}} \pm 0.08_{\text{syst}}) 10^{-6}$$

Last measurement: BNL E787(1991):

31 candidates, 5 BG events:

$$BR = (1.10 \pm 0.32) 10^{-6}$$




Conclusions

- an accurate study of low energy QCD based on highest statistics of K_{e4} is achieved in the NA48/2 experiment:
 - 5 FF's are measured with high precision
 - the measured pi-pi scattering lengths are in very good agreement with the "cusp result" & ChPT prediction
- the first measurement of INT term in $K^{\pm} \rightarrow \pi^{\pm} \pi^0 \gamma$ decay & CPV test are performed
- the $\text{Br}(K^{\pm} \rightarrow \pi^{\pm} \gamma \gamma)$ is measured with the highest precision


Spare

NA48/2 beam line



2-3M K/spill ($\pi/K \sim 10$),
 π decay products stay in pipe.
Flux ratio: $K^+/K^- \approx 1.8$

Simultaneous K^+ and K^- beams:
large **charge symmetrization** of
experimental conditions



Front-end achromat

Quadrupole quadruplet

• Cleaning
• Beam spectrometer (resolution 0.7%)

Beams coincide within ~ 1 mm
all along 114m decay volume


• Momentum selection

• Focusing
• μ sweeping

10 cm

Ke4 phase shift measurement

isospin correction (10-15 mrad) to all points are applied - downward shift errors — combined stat. and syst.; fit to NA48/2 data alone



Systematic errors:

- bin to bin uncorrelated:
 - fitting procedure*
 - trigger efficiency*
 - acceptance control*
 - background shape*
 - electron identification*
 - radiative correction*
- bin to bin correlated:
 - background level*
 - isospin corrections*

$$\mathbf{K}^{\pm} \rightarrow \pi^0 e^{\pm} \nu \gamma$$



260k decays

The measurements of:

➤ BR

➤ CPV test

Ke3 γ : Theory (ChPT)



$$\text{BR}(\text{Ke}3\gamma/\text{Ke}3) = (6.40 \pm 0.08) \cdot 10^{-3} \quad (E^* > 30 \text{ MeV}, \theta_{e\gamma}^* > 20^\circ)$$

Computed in *Eur. Phys. Jour. C* 50 (2007)

$$\xi = \frac{\vec{p}_\gamma \cdot (\vec{p}_\pi \times \vec{p}_e)}{M_K^3}$$

$N_{+(-)}$ = number of events with positive (negative) ξ

$$A_\xi = \frac{N_+ - N_-}{N_+ + N_-}$$

$(A_\xi^+ + A_\xi^-)$ directly probes CPV