



Contribution ID: 134

Type: Parallel Session Talk

Study light scalar mesons from heavy quark decays

Friday, 23 July 2010 12:15 (13 minutes)

It is a difficult task to probe internal structures of the scalar mesons below or near 1GeV. In the SU(3) symmetry limit, the semileptonic $D^+ \rightarrow Sl^+ \nu$ and $B^+ \rightarrow Sl \text{ nubar}$ decays, with $S=a_0, f_0$ and σ , are found to obey very different sum rules in the two scenarios for scalar mesons. Thus it can uniquely distinguish the $q\bar{q}$ and the tetraquark descriptions for light scalar mesons model-independently. This also applies to the $B^0 \rightarrow J/\psi(\eta_c) S$ decays. The SU(3) symmetry breaking effect is found to be under control, which will not spoil our method. The branching fractions of the $D^+ \rightarrow Sl^+ \nu$, $B^+ \rightarrow Sl \text{ nubar}$ and $B^0 \rightarrow J/\psi(\eta_c) S$ decays roughly have the order 10^{-4} , 10^{-5} and 10^{-6} , respectively. The ongoing BES-III and the forthcoming Super B experiments are able to measure these channels and accordingly to provide the detailed information of scalar meson inner structure.

Primary author: Prof. LU, Cai-Dian (IHEP,Beijing)

Co-author: Dr WANG, Wei (INFN, Bari,)

Presenter: Prof. LU, Cai-Dian (IHEP,Beijing)

Session Classification: 04 - Hadronic Structure, Parton Distributions, soft QCD, Spectroscopy

Track Classification: 04 - Hadronic Structure, Parton Distributions, soft QCD, Spectroscopy