


# Measurement of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ Decay at Fermilab

## $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ in the Standard Model

The  $K \rightarrow \pi \nu \bar{\nu}$  decays are the most precisely calculated FCNC decays.




- A single effective operator  $(\bar{s}_L \gamma^\mu d_L)(\bar{\nu}_L \gamma_\mu \nu_L)$
- Dominated by top quark (charm significant, but controlled)
- Hadronic matrix element shared with  $K \rightarrow \pi e \nu$
- Largest uncertainty from CKM elements (which will improve)


$$B_{SM}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (8.5 \pm 0.7) \times 10^{-11}$$

Brod and Gorbahn, PRD 78, 034006(2008)

- Remains clean in New Physics models




Measurement of  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  at a level exceeding the highly precise SM prediction could signal the presence of new physics at high mass scales. Observation of  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  with a  $10^{-10}$  branching ratio is experimentally challenging. The signature is charged kaon followed by a charged pion with no other observed particles. Backgrounds from  $K^+ \rightarrow \pi^+ \pi^0$  and  $K^+ \rightarrow \mu^+ \nu$  involve branching ratios ten orders of magnitude larger. The experimental strategy is to prove that candidate events have a low probability of being due to background. Successful  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  detection requires powerful  $\pi^+$  particle identification, high-efficiency,  $4\pi$  sr photon veto capability and efficient  $K^+$  identification to eliminate beam-related background.



Over 2 decades, BNL experiments E787 and E949 refined the technique for  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  detection using stopped kaons over 3 generations of experiments to observe 7  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  candidates resulting in  $B(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (1.73^{+1.15}_{-1.05}) \times 10^{-10}$ .

## General MSSM with R-parity

Effect of a  $\pm 5\%$  measurement at a hypothetical non-SM BF




Buras et al, NP B714, 103(2005)

Points from a scan of MSSM parameters that satisfy experimental constraints except  $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$

E949 already provides a significant constraint.


New Physics models with generic flavor structure typically induce large effects in these decays.

Measured range in plastic scintillator vs kinetic energy for all  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  candidates in E787 & E949. The solid and dashed lines indicate the signal regions. The grey points are from signal MC. The cluster of points near 108 MeV is due to  $K^+ \rightarrow \pi^+ \pi^0$  events surviving the photon veto.



PRL 93, 031801(2004). PRD 77, 052003(2008). PRL 101, 191802(2008). PRD 79, 092004(2009).

## P996 DETECTOR




Proposed P996 detector.

A 550 MeV/c  $K^+$  beam is stopped in a highly segmented active target.  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  decays are observed with a precision, low-mass central drift chamber surrounded by segmented scintillator detectors to measure pion range, energy and the  $\pi^+ \mu^+ e^-$  decay sequence and enclosed by an efficient  $4\pi$  sr EM calorimeter for vetoing events accompanied by photons.

The fourth generation experiment, P996, would detect about 200  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  decays per year by exploiting incremental improvements over E949, a high duty factor utilizing the Tevatron as a "Stretcher" and longer running time per year.

## P996 sensitivity



A precision of  $<5\%$  could be achieved in 3-5 years if the branching ratio is consistent with the SM expectation.

Member institutions of the P996 collaboration are

Arizona State University(USA), Brookhaven National Laboratory(USA), Fermilab(USA), Institute for Nuclear Research(Russia), Istituto Nazionale di Fisica Nucleare, Pisa (Italy), JINR, Dubna (Russia) TRIUMF(Canada), University of British Columbia(Canada), University of Texas at Austin(USA), University of Illinois, Urbana(USA), University of Northern British Columbia(Canada), Universidad Autonoma de San Luis Potosi(Mexico), Tsinghua University, Beijing(China)