

Z'_{B-L} discovery potential at the LHC for $\sqrt{s} = 7$ TeV

Lorenzo Basso (UoS and RAL)

In collaboration with: S. Moretti, A. Belyaev, G.M. Pruna,
C.H. Shepherd-Themistocleous

LB, A. Belyaev, S. Moretti, C.H. Shepherd-Themistocleous: Phys. Rev. D **80**, 055030 (2009) [hep-ph:0812.4313]
LB, A. Belyaev, S. Moretti, G.M. Pruna: JHEP **0910**, 006 (2009) [hep-ph:0903.4777]
LB, A. Belyaev, S. Moretti, G.M. Pruna, C.H. Shepherd-Themistocleous: hep-ph:1002.3586

Outline

1) The $B - L$ model

- Motivations
- Particle content

2) Phenomenology (at the LHC)

- Z' properties: decays and width
- Analysis details
- LHC @ $\sqrt{s} = 7$ TeV
- Tevatron comparison
- Exclusions

3) Appendix: Heavy neutrino phenomenology (at the LHC)

- Decays and width
- Tri-lepton signature

4) Conclusions

A triply minimal extension

- **Motivations**

- ▷ ν_R naturally included
- ▷ $U(1)_{B-L}$ (accidental) global symmetry in the SM
- ▷ interesting phenomenology
- ▷ part of a bigger picture (GUT, baryogenesis)

- **Gauge sector**

$$SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$$

- **Fermion sector**

One extra fermion per generation: ν_R
(Required by anomaly cancellation)

- **Scalar sector**

One extra SM-singlet scalar: χ
($U(1)_{B-L}$ symmetry breaking)

The model: triply-minimal extension

A $U(1)$ extension of the SM

$$SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$$

New states:

- A scalar (χ , SM-singlet)

$$V = \dots + \lambda_1 (H^\dagger H)^2 + \lambda_2 |\chi|^4 + \lambda_3 H^\dagger H |\chi|^2$$

- 3 RH neutrinos: $\nu_R \xrightarrow{\text{see-saw}} \nu_h$ ($\mathcal{O}(100)$ GeV)
(anomaly cancellation)

$$\mathcal{L}_Y = \dots - y^\nu \bar{l}_L \nu_R \tilde{H} - y^M \overline{(\nu_R)^c} \nu_R \chi + \text{H.c.}$$

In certain regions of the parameter space, they both can be *long-lived* particles

ψ	$SU(3)_C$	$SU(2)_L$	Y	$B-L$
q_L	3	2	$\frac{1}{6}$	$\frac{1}{3}$
u_R	3	1	$\frac{2}{3}$	$\frac{1}{3}$
d_R	3	1	$-\frac{1}{3}$	$\frac{1}{3}$
l_L	1	2	$-\frac{1}{2}$	-1
e_R	1	1	-1	-1
ν_R	1	1	0	-1

ψ	$SU(3)_C$	$SU(2)_L$	Y	$B-L$
H	1	2	$\frac{1}{2}$	0
χ	1	1	0	2

Interactions and spectrum

Covariant derivative (in a suitable basis):

$$D_\mu \Psi_i = \partial_\mu \Psi_i + i [g_1 Y_i B_\mu + (Y_i \tilde{g} + (B - L)_i g'_1) B'_\mu] \Psi_i$$

$Z - Z'$ mixing:

$$\begin{pmatrix} B^\mu \\ W_3^\mu \\ B'^\mu \end{pmatrix} = \left(\begin{array}{cc|c} \cos \vartheta_w & -\sin \vartheta_w \cos \vartheta' & \sin \vartheta_w \sin \vartheta' \\ \sin \vartheta_w & \cos \vartheta_w \cos \vartheta' & -\cos \vartheta_w \sin \vartheta' \\ 0 & \sin \vartheta' & \cos \vartheta' \end{array} \right) \begin{pmatrix} A^\mu \\ Z^\mu \\ Z'_{B-L}{}^\mu \end{pmatrix}$$



$$\tilde{g} = 0 \longrightarrow \vartheta' = 0$$

No (tree level) $Z - Z'$ mixing in the pure $B - L$ model

Z' phenomenology

$m_{\nu h} = 250 \text{ GeV}$

1 TeV



$$\sum_k BR(Z'_{B-L} \rightarrow l_k \bar{l}_k) \sim \frac{3}{4}$$

$$\sum_k BR(Z'_{B-L} \rightarrow q_k \bar{q}_k) \sim \frac{1}{4}$$

- Dominantly coupled to *leptons*
- $Z' \rightarrow \nu_h \nu_h$ up to $\sim 20\%$

- $g'_1 < 0.5$ from RGE analysis
- Γ up to hundreds of GeV

Depending on $m_{\nu h}$ and $M_{Z'}$:

$$Z' \rightarrow l^+ l^- \simeq 12.5\% \div 15.5\%$$

$$Z' \rightarrow q \bar{q} \simeq 4\% \div 5\%$$

In this work: $m_{\nu h} = 200 \text{ GeV}$

Aim of our work

The $B - L$ symmetry breaking takes place at the TeV scale. Its rich phenomenology (Z' , Higgs(es) and heavy neutrinos signatures) can be observed at TeV machines.

We focus our analysis on the Z' -sector at the LHC at its current stage, $\sqrt{s} = 7$ TeV.

We consider the $pp(\bar{p}) \rightarrow l^+l^-$ ($l = e, \mu$) channel as a representative process to study new signatures of the Z'_{B-L} vector boson.

The $B - L$ model is not a usual benchmark for data analyses nor colliders reach studies \rightarrow *systematic parton level analysis, with k -factors*.
(no ISR, photon conversions, fakes, muon reconstructions, ...)

Differences wrt the other more common Z' models:

- no axial coupling: $(B - L)^R = (B - L)^L$ hence $g_{Z'}^A = \frac{g_{Z'}^R - g_{Z'}^L}{2} = 0$,
- free value of the coupling g'_1 ,
- new coupled matter (ν_h) \rightarrow not-fixed BRs.

Analysis details

Analysis with CalcHEP, model implemented via LanHEP.

We have assumed standard acceptance cuts at LHC and Tevatron (the same for electrons and muons):

$$\begin{aligned} \text{LHC :} \quad p_T &> 10 \text{ GeV}, & |\eta| < 2.5, \\ \text{Tevatron :} \quad p_T &> 18 \text{ GeV}, & |\eta| < 1, \end{aligned}$$

and a cut on the invariant mass (**early** vs **improved** scenario):

$$\begin{aligned} e^\pm @ \text{LHC:} \quad \frac{|M_{ee} - M_{Z'}|}{\text{GeV}} &< \max \left(\frac{\Gamma_{Z'}}{2}, \left(\begin{array}{c} 0.02 \\ 0.005 \end{array} \right) \frac{M_{Z'}}{\text{GeV}} \right), \\ \mu^\pm @ \text{LHC:} \quad \frac{|M_{\mu\mu} - M_{Z'}|}{\text{GeV}} &< \max \left(\frac{\Gamma_{Z'}}{2}, \left(\begin{array}{c} 0.08 \\ 0.04 \end{array} \right) \frac{M_{Z'}}{\text{GeV}} \right), \end{aligned}$$

Analysis details

Analysis with CalcHEP, model implemented via LanHEP.

We have assumed standard acceptance cuts at LHC and Tevatron (the same for electrons and muons):

$$\begin{aligned} \text{LHC :} \quad p_T &> 10 \text{ GeV}, & |\eta| < 2.5, \\ \text{Tevatron :} \quad p_T &> 18 \text{ GeV}, & |\eta| < 1, \end{aligned}$$

and a cut on the invariant mass (**early** vs **improved** scenario):

$$e^\pm @ \text{LHC: } \frac{|M_{ee} - M_{Z'}|}{\text{GeV}} < \max \left(\frac{\Gamma_{Z'}}{2}, \left(\begin{matrix} 0.02 \\ 0.005 \end{matrix} \right) \frac{M_{Z'}}{\text{GeV}} \right),$$

$$\mu^\pm @ \text{LHC: } \frac{|M_{\mu\mu} - M_{Z'}|}{\text{GeV}} < \max \left(\frac{\Gamma_{Z'}}{2}, \left(\begin{matrix} 0.08 \\ 0.04 \end{matrix} \right) \frac{M_{Z'}}{\text{GeV}} \right),$$

$$e^\pm @ \text{Tevatron: } \frac{|M_{ee} - M_{Z'}|}{\text{GeV}} < \max \left(\frac{\Gamma_{Z'}}{2}, \left(0.135 \sqrt{\frac{M_{Z'}}{\text{GeV}}} + 0.02 \frac{M_{Z'}}{\text{GeV}} \right) \right),$$

$$\mu^\pm @ \text{Tevatron: } \frac{|M_{\mu\mu} - M_{Z'}|}{\text{GeV}} < \max \left(\frac{\Gamma_{Z'}}{2}, 0.0005 \left(\frac{M_{Z'}}{2 \text{ GeV}} \right)^2 \right).$$



Z' Discovery potentials for $\sqrt{s} = 7$ TeV

Significance contour levels plotted against g'_1 and $M_{Z'}$

Electrons

↓ Tevatron and LEP bounds ↓


Muons




Z' Discovery potentials for $\sqrt{s} = 7$ TeV

Luminosity vs $M_{Z'}$ (for fixed values of g_1') at the LHC

Electrons



Muons




Z' Discovery potentials at the Tevatron

Luminosity vs $M_{Z'}$ (for fixed values of g'_1)

Electrons

Electrons Tevatron ($\sqrt{s}=1.96$ TeV)


— $5\sigma, g'_1=0.08$ — $5\sigma, g'_1=0.1$
- - $3\sigma, g'_1=0.08$ - - $3\sigma, g'_1=0.1$



Muons


Muons Tevatron ($\sqrt{s}=1.96$ TeV)

— $5\sigma, g'_1=0.08$ — $5\sigma, g'_1=0.1$
- - $3\sigma, g'_1=0.08$ - - $3\sigma, g'_1=0.1$



Z' exclusions @ 7 TeV

Significance contour levels plotted against g'_1 and $M_{Z'}$



Z' exclusions @ 7 TeV (II)

Luminosity vs $M_{Z'}$ (for fixed values of g'_1)

LHC ($\sqrt{s}=7$ TeV), 95% C.L.

Electrons

— $g'_1=0.158$

— $g'_1=0.1$


— $g'_1=0.05$

Muons

- - - $g'_1=0.158$

- - - $g'_1=0.1$

- - - $g'_1=0.05$



Tevatron, 95% C.L.

Electrons


— $g'_1 = 0.1$

— $g'_1 = 0.05$



Muons

- - - $g'_1 = 0.1$

- - - $g'_1 = 0.05$



Appendix: ν_h phenomenology



$$\tan 2\alpha_\nu = -2\sqrt{\frac{m_{\nu l}}{m_{\nu h}}}$$

Feynman diagram showing the decay of ν_h into a lepton l and a W boson. The vertex is labeled with the expression:


$$= \frac{\sqrt{2}e}{4 \sin \vartheta_W} \sin \alpha_\nu$$

- $\Gamma = \Gamma(m_{\nu l}/m_{\nu h})$
- ν_h can be a long-lived particle
- **DISPLACED VERTICES** (e.g., from a high energetic and isolated lepton)

ν_h @ LHC: $BR(Z' \rightarrow 3l + 2j + \cancel{P}_T (1\nu), l = e, \mu)$ up to 2.5%

$$m_T^2 = \left(\sqrt{M_{vis}^2 + P_{T,vis}^2} + |\cancel{P}_T| \right)^2 - \left(\vec{P}_{T,vis} + \vec{\cancel{P}}_T \right)^2$$

V. Barger et al, Phys. Rev. D **36** (1987) 295



$M_{Z'} = 1.5 \text{ TeV}, g'_1 = 0.2: \sigma(pp \rightarrow Z') = 0.3 \text{ pb}$
 $M_{\nu_h} = 200 \text{ GeV}, \mathcal{L} = 100 \text{ fb}^{-1}, \text{bin} = 20 \text{ GeV}$

Backgrounds:

$WZjj$ associated production ($\sigma_{3l} = 246.7 \text{ fb}, l = e, \mu, \tau, \text{w. cuts}$)

$t\bar{t}$ pair production ($\sigma_{2l} = 29.6 \text{ pb}, l = e, \mu$) (3rd lep. from b-quark)



$t\bar{t}l\nu$ associated production ($\sigma_{3l} = 8.6 \text{ fb}, l = e, \mu, \tau$)

Cuts:

Kinematics, angular acceptance and isolation

W rec. from jets: $|M_{jj} - 80 \text{ GeV}| < 20 \text{ GeV}$

Z' rec.: $|M_{3l,2j}^T - 1500 \text{ GeV}| < 250 \text{ GeV}$



Conclusions

- ✓ Simple SM extension at TeV scale, RH-neutrinos
- ✓ motivated by high-scale physics (GUT, baryogenesis)
- ✓ pure $B - L$ model, no $Z - Z'$ mixing, only vectorial coupling

- Discovery power limited by existing experimental constraints
good prospects at LHC @ $\sqrt{s} = 7$ TeV
- Better sensitivity to electrons (also, width measure possible)


Tevatron@10 fb⁻¹ stil competitive with LHC@7 TeV:

- LHC requires 1 fb⁻¹ for same sensitivity to small couplings
- but it is not limited by kinematics: 5σ observation at LHC@7 TeV up to $M_{Z'} = 1.25(1.20)$ TeV with electrons(muons)
- Otherwise, 95% C.L. exclusions

What's next: look for ν_h

Backup slides

Line shape



Z' width measure possible with electrons

Validation

Comparison of our simulation (CalcHEP) wrt $D\bar{D}$ analysis of expected backgrounds ¹ at 3.6 fb^{-1} of integrated luminosity. From Table II in

<http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/NP/N66/>.


The compatibility is, as usual, defined as:

$$\frac{Ev - Ev_{D\bar{D}}}{\sigma_{D\bar{D}}}$$

$M_{Z'}$ (GeV)	Mass Window Lower Limit (GeV)	$D\bar{D}$ analysis		Our simulation	
		Expected Bkg Events	Signal Acceptance	Expected Bkg Events	Compatibility Level
400	353	22.4 ± 0.7	0.172	23.2	1.1
500	445	7.92 ± 0.22	0.188	8.23	1.4
600	536	2.93 ± 0.07	0.199	3.00	1.0
700	626	1.052 ± 0.025	0.207	1.110	2.3
750	673	0.631 ± 0.016	0.209	0.653	1.4
800	718	0.384 ± 0.010	0.211	0.391	0.7
850	762	0.222 ± 0.006	0.212	0.235	2.2
900	810	0.134 ± 0.004	0.216	0.135	0.4
950	858	0.0701 ± 0.0023	0.214	0.0750	2.1
1000	902	0.0410 ± 0.0015	0.216	0.0440	2.0

¹The $D\bar{D}$ simulation is modelled using the PYTHIA Monte Carlo event generator, with CTEQ6L1 PDFs, and then processed through the standard $D\bar{D}$ detector simulation based on GEANT3.

CMS resolution vs Z' width



Peak dominated by experimental resolution if $g'_1 \lesssim 0.4(0.2)$



Z' Discovery potentials for $\sqrt{s} = 14$ TeV

Significance contour levels plotted against g'_1 and $M_{Z'}$, improved resolutions.

Electrons

↓ Tevatron and LEP bounds ↓


Muons




Z' Discovery potentials for $\sqrt{s} = 14$ TeV

Luminosity vs $M_{Z'}$ (for fixed values of g'_1)

Electrons




Muons




Z' exclusions @14 TeV

Significance contour levels plotted against g'_1 and $M_{Z'}$



Luminosity vs $M_{Z'}$
(for fixed values of g'_1)



Z' experimental limit


LEP bound:

G. Cacciapaglia *et al*, Phys. Rev. D **74** (2006) 033011

$$\frac{M_{Z'}}{g'_1} \geq 7 \text{ TeV}$$

Tevatron (comparing our prediction to experimental data):

Muons T. Aaltonen *et al*. [CDF], Phys. Rev. Lett. **102**, 091805 (2009):




$p\bar{p} \rightarrow \mu^+ \mu^-$	
g'_1	$M_{Z'} \text{ (GeV)}$
0.06	600
0.1	750
0.123	800
0.2	900
0.3	1000
0.5	1195

Z' experimental limit: continue

Tevatron (comparing our prediction to experimental data):

Electrons T. Aaltonen *et al.* [CDF], Phys. Rev. Lett. **102**, 031801 (2009):



$p\bar{p} \rightarrow e^+e^-$	
g'_1	$M_{Z'} \text{ (GeV)}$
0.042	600
0.086	700
0.115	800
0.19	900
0.3	1000