



Contribution ID: 1092

Type: Parallel Session Talk

CP violation and mixing in charm decays at Belle

Friday, 23 July 2010 17:15 (13 minutes)

We have searched for CP violation of charmed mesons in the decays $D \rightarrow K_0^* P$, where D denotes D^0, D^+ and D_s^+ , and P denotes the pseudo-scalar mesons π^+, K^+, π^0, η and η' . No evidence of CP violation in these decays is observed. We have measured the CP asymmetry difference between the Cabibbo suppressed decay $D^+ \rightarrow \phi \pi^+$ and the Cabibbo favored $D_s^+ \rightarrow \phi \pi^+$ decays. The measured asymmetry is corrected for the residual asymmetry due to detector effects, and the contributions of both CP and forward-backward asymmetries are determined. We also present a measurement of D^0 -anti- D^0 mixing parameters in three-body D^0 decays using a time-dependent Dalitz plot analysis. These results are obtained on a large data sample collected at and near the $\Upsilon(4S)$ resonance with the Belle detector operating at the KEKB asymmetric-energy $e^+ e^-$ collider.

Primary author: BELLE, Collaboration (KEK)**Presenter:** KO, Byeong Rok**Session Classification:** 06 - CP violation, CKM and Rare Decays**Track Classification:** 06 - CP violation, CKM and Rare Decays