



Contribution ID: 984

Type: Parallel Session Talk

## Semileptonic D Meson Decays from CLEO-c

*Saturday, 24 July 2010 09:15 (15 minutes)*

Using the full CLEO-c  $D^0D^0\bar{0}$ ,  $D^+D^-$ , and  $D_s\bar{D}_s$  data samples, we have made precision measurements of many D meson semileptonic branching fractions and form factors. The results for the form factors of exclusive  $D^+$  and  $D^0$  semileptonic decays to K and  $\pi$  mesons agree well with recent Lattice QCD calculations. Using a non-parametric technique, we measure the form factor for  $D^{+-}\rightarrow K^0\bar{0} e^+ \nu_e$  decay and also the mass-suppressed form factor in  $D^{+-}\rightarrow K^0\bar{0} \mu^+ \nu_\mu$  decay. We report new results for the Cabibbo-suppressed semileptonic decays to the vector mesons,  $D^0\rightarrow\rho^- e^+ \nu_e$ ,  $D^{+-}\rightarrow\rho^0 e^+ \nu_e$ , and  $D^{+-}\rightarrow\omega e^+ \nu_e$ , as well as to the scalars  $\eta$  and  $\eta'$ . We also report measurements of exclusive  $D_s$  semileptonic decays to a variety of final states. Finally, we report precision measurements of inclusive  $D^0$ ,  $D^+$ , and  $D_s$  semileptonic decays.

**Primary author:** CASSEL, David (Cornell University)

**Presenter:** CASSEL, David (Cornell University)

**Session Classification:** 05 - Heavy Quarks Properties (experiment and theory)

**Track Classification:** 05 - Heavy Quarks Properties (experiment and theory)