

# Is the dark matter particle its own anti-particle?

**Z. Chacko**

**University of Maryland, College Park**

# **Introduction**

**It is now well established from astrophysical and cosmological observations that about 80% of the matter in the universe is made up of non-luminous, non-baryonic dark matter.**



**However, the nature of the particles of which dark matter is composed remains a mystery.**

**Stable particles with masses of order the weak scale that have weak scale cross sections with visible matter naturally have the right relic abundance to explain observations - 'the WIMP miracle'.**

**Since the WIMP dark matter framework is so robust, it naturally arises or can be accommodated in many different scenarios.**

**To what extent is it possible to extract from experiment information about the nature of the dark matter particle independent of any specific model?**

**Clearly, detailed predictions will not be possible.**

**However, it may be possible to answer several of the most important questions about the dark matter particle.**

**• What is the mass of the dark matter particle?**

**• What is its spin?**

**• Is the dark matter particle its own anti-particle?**

**• What are its SM quantum numbers?**




Let us formulate the problem.

According to the *CPT* theorem, for every particle there is an anti-particle with the same mass and spin, but opposite charge(s).

Since dark matter is electrically neutral, it is possible that the dark matter particle is its own anti-particle.

More precisely,



Can dark matter experiments address this question?

The logic of this talk is as follows.

I will establish a very close correlation between theories where the WIMP-nucleon cross section is dominated by spin-dependent interactions, and theories where the dark matter particle is its own anti-particle.

$$\sigma_{\text{SD}} \gg \sigma_{\text{SI}} \quad \longrightarrow \quad \chi = \chi^c$$

However

$$\chi = \chi^c \quad \not\longrightarrow \quad \sigma_{\text{SD}} \gg \sigma_{\text{SI}}$$

The experiment that is most sensitive to spin-dependent dark matter is the IceCube neutrino telescope located at the South Pole. A signal could help establish the dark matter particle is its own anti-particle!

The problem is that IceCube is also sensitive to spin-independent interactions. Not possible to distinguish the origin of the signal.

However, I will then show that limits from direct detection expts. can be used to place a model-independent upper bound on the event rate from spin-independent interactions, closing the loophole.

# **Spin-dependent dark matter candidates**

We wish to classify theories which can naturally lead to primarily spin-dependent interactions with matter.

Cosmological limits constrain WIMP dark matter to be neutral under color and electromagnetism.

We will limit to theories where WIMP-nucleon scattering is elastic and arises from an effective operator generated by a tree diagram at parton level → only WIMP-quark operators need be considered.

In this class of theories WIMP-nucleon scattering is generated by operators of the general form

$$\frac{[\text{dark matter bilinear}] [\text{quark bilinear}]}{M^n}$$

Here  $M$  is the mass of the particle mediating the interaction, and  $n$  is less than or equal to 2. In general the bilinears include derivatives.



The cross section depends on the matrix element of the quark bilinear between nuclear states. Parity symmetry allows us to distinguish spin-dependent terms.

Let  $s$  represent the spin of the nucleus and  $v$  the relative velocity of the WIMP-nucleus system. While  $s$  is a pseudo-vector,  $v$  is a vector. Other parameters, such as reduced mass and charge(s) are scalars.

scalar

$$\bar{q} q$$

pseudo-scalar

$$\bar{q} \gamma^5 q \sim \vec{s} \cdot \vec{v}$$

vector

$$\bar{q} \gamma^\mu q \begin{cases} \rightarrow \bar{q} \gamma^0 q \\ \rightarrow \bar{q} \gamma^i q \sim v^i \end{cases}$$

pseudo-vector

$$\bar{q} \gamma^\mu \gamma^5 q \begin{cases} \rightarrow \bar{q} \gamma^0 \gamma^5 q \sim \vec{s} \cdot \vec{v} \\ \rightarrow \bar{q} \gamma^i \gamma^5 q \sim s^i \end{cases}$$

tensor

$$\bar{q} \sigma^{\mu\nu} q \begin{cases} \rightarrow \bar{q} \sigma^{0i} q \sim v^i \\ \rightarrow \bar{q} \sigma^{ij} q \sim \epsilon^{ijk} s^k \end{cases}$$

The velocity dependent terms can be neglected, since  $v \sim 10^{-3}$

Of the remaining terms,  $\bar{q}q$  and  $\bar{q}\sigma^{\mu\nu}q$

are only generated by effects that break chiral symmetry, and can naturally be small.

Then, for spin-dependent interactions to dominate terms involving

$\bar{q}\gamma^\mu\gamma^5q$  must be present in the theory, while terms involving the

operator  $\bar{q}\gamma^\mu q$  must be absent.

What are the theories where this can happen naturally?

The theories where spin-dependent interactions can naturally dominate involve either Majorana fermions or real vector bosons.  
→ The dark matter particle has spin and is its own anti-particle.

## Scalar WIMPs

Scalars always lead to spin-independent interactions.

## Fermionic WIMPs

There are two operators that can lead to scattering.

$$\bar{\chi} \gamma_{\mu} \gamma^5 \chi \bar{q} \gamma^{\mu} \gamma^5 q \longrightarrow \text{SD}$$

$$\bar{\chi} \gamma_{\mu} \chi \bar{q} \gamma^{\mu} q \longrightarrow \text{SI}$$

For a Majorana fermion the second operator vanishes  $\rightarrow$  scattering is spin-dependent. For Dirac case (in general) both operators contribute.

## Vector boson WIMPs

For real vector bosons only one operator contributes in chiral limit.

$$\epsilon_{\mu\nu\lambda\sigma} \partial^{\mu} B^{\nu} B^{\lambda} \bar{q} \gamma^{\sigma} \gamma^5 q \longrightarrow \text{SD}$$

For complex vector bosons an additional operator contributes.

$$\partial_{\mu} B^{*}_{\nu} B^{\nu} \bar{q} \gamma^{\mu} q \longrightarrow \text{SI}$$

Dark Matter	Mediator	Process	Scattering
Scalar	$Z, Z'$		SI
	$h$		SI
	$Q$		SI
Dirac Fermion	$Z, Z'$		SI, SD <sup>†</sup>
	$h$		SI
	$X$		SI, SD
	$\Phi$		SI, SD
Majorana Fermion	$Z, Z'$		SD
	$h$		SI
	$X$		SD in chiral limit
	$\Phi$		SD in chiral limit
Real Vector	$h$		SI
	$Q$		SD in chiral limit
Complex Vector	$Z, Z'$		SI
	$h$		SI
	$Q$		SI, SD


Table 1: A summary of results for WIMP-nucleon scattering, for each dark matter candidate and mediator [36]. In the Feynman diagrams, scalars are represented by dashed lines, fermions by solid lines and vector bosons by wavy lines. Of the mediators,  $h$ ,  $Z'$  and the SM  $Z$  are neutral under both electromagnetism and color, while  $X$ ,  $\Phi$  and  $Q$  transform as triplets under color and carry electric charge.

<sup>†</sup>Can be primarily SD for specific choices of  $Z'$  charges


# **Limits on dark matter event rates in IceCube**

**Neutrino telescopes are searching for neutrinos arising from dark matter annihilation in the sun (and earth).**

**Dark matter particles collide with nuclei in the sun, thereby losing energy. As a consequence they become gravitationally bound.**




**After subsequent scatterings, accumulate in the core of the sun. Can then pair annihilate, giving rise to neutrinos.**



**These neutrinos can give rise to a signal in neutrino telescopes.**

The limits from direct detection experiments can be used to place a model-independent bound on the dark matter event rate in IceCube.


How? Capture in the sun occurs through WIMP-nucleon scattering, which is bounded by direct detection experiments. The direct detection limit translates into a bound on the WIMP capture rate.



The annihilation rate can at most be equal to half the capture rate. The limit on the capture rate is then also a limit on the annihilation rate.

For a fixed dark matter mass and annihilation into the most neutrino rich final state, leads to an upper limit on the total neutrino event rate.

For spin-independent interactions, this limit is comparable to current IceCube bound.



**Spin-independent interactions could still be responsible for generating a signal at IceCube close to the current experimental bound, if annihilation is directly to neutrino rich final states. But . . .**

**The limits from direct detection on spin-independent interactions are expected to improve by two orders of magnitude in the near future.**

**At that point, any observed signal at IceCube must be arising from spin-dependent interactions → a strong hint that the dark matter particle is its own anti-particle.**



# Conclusions

**The dark matter candidates that naturally tend to have primarily spin-dependent interactions with matter are Majorana fermions and real vector bosons, so that the dark matter particle is its own anti-particle.**

**IceCube is currently sensitive to both spin-independent as well as spin-dependent dark matter candidates.**

**If the direct detection bounds continue to improve, the case of spin-independent dark matter will soon go out of reach of IceCube.**

**In such a scenario, a signal at IceCube would constitute a strong hint that the dark matter particle has spin, and is its own anti-particle.**

**The region of parameter space where IceCube can expect a signal from spin-dependent dark matter is kinematically accessible to LHC.**

# **IceCube and the LHC**

If the interactions of dark matter with nuclei are spin-dependent, the natural candidates are Majorana fermions and real vector bosons.

If the dark matter candidate is a Majorana fermion, spin-dependent WIMP-nucleon scattering can arise through any of:

• t-channel vector exchange

SM Z  
Z'

WIMP carries weak charge

new Z' in the theory

• s- and u-channel vector exchange

mediator carries SM color

• s- and u-channel scalar exchange

mediator carries SM color


If the dark matter candidate is a real vector, only one possibility:

• s- and u-channel fermion exchange

mediator carries SM color

Either new Z', or new states with SM charges → promising for LHC!

The figure shows the range of mediator masses that leads to a signal at IceCube with 5 years of data. Promising for the LHC.



The case with a new  $Z'$  is disfavored by precision electroweak limits.

The new colored particles are kinematically accessible to the LHC.

For Majorana fermion dark matter charged under the SM Z, some part of the parameter space is kinematically accessible to the LHC.

Direct detection experiments can expect to see a small but definite spin-independent signal.

