



2HDMC: two Higgs doublet model calculator

Johan Rathsman, Lund and Uppsala University

Charged2010, Sätra Brunn, 2010-09-29

- ➊ General two Higgs doublet models (2HDM)
- ➋ Theoretical constraints
- ➌ Yukawa sector
- ➍ Experimental constraints
- ➎ Usage/setting parameters/input-output



2HDMC

Johan Rathsman

2HDM

2HDM potential

EWSB

Symmetries

TH constraints

Yukawa sector

EX constraints

Usage

Example

Reference

D. Eriksson, JR and O. Stål, Comp. Phys. Comm. **181** (2010) 189; 833
<http://www.isv.uu.se/theep/MC/2HDMC>



General two Higgs doublet model potential

- Two complex $SU(2)_L$ doublets with hypercharge $Y=1$: Φ_1, Φ_2
- Invariance under global $SU(2)$: $\Phi_a \rightarrow U_{ab} \Phi_b$

General potential

$$\begin{aligned}\mathcal{V} = & m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 - \left[m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right] + \frac{1}{2} \lambda_1 \left(\Phi_1^\dagger \Phi_1 \right)^2 \\ & + \frac{1}{2} \lambda_2 \left(\Phi_2^\dagger \Phi_2 \right)^2 + \lambda_3 \left(\Phi_1^\dagger \Phi_1 \right) \left(\Phi_2^\dagger \Phi_2 \right) + \lambda_4 \left(\Phi_1^\dagger \Phi_2 \right) \left(\Phi_2^\dagger \Phi_1 \right) \\ & + \left\{ \frac{1}{2} \lambda_5 \left(\Phi_1^\dagger \Phi_2 \right)^2 + \left[\lambda_6 \left(\Phi_1^\dagger \Phi_1 \right) + \lambda_7 \left(\Phi_2^\dagger \Phi_2 \right) \right] \left(\Phi_1^\dagger \Phi_2 \right) + \text{h.c.} \right\}\end{aligned}$$

- Potential real $\Rightarrow m_{11}^2, m_{22}^2, \lambda_{1-4}$ real m_{12}^2, λ_{5-7} complex
- No explicit CP-violation $\Rightarrow m_{12}^2, \lambda_{5-7}$ real



Electroweak symmetry breaking

- EW symmetry broken by non-zero vev of Φ_1 and/or Φ_2
- Minimization conditions $\Rightarrow m_{11}^2, m_{22}^2$ traded for $v_1 = v \cos \beta, v_2 = v e^{i\xi} \sin \beta$ with $v = (\sqrt{2} G_F)^{-1/2} \approx 246$ GeV
- No spontaneous CP-violation $\Rightarrow \xi = 0$

$$\Phi_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}(G^+ \cos \beta - H^+ \sin \beta) \\ v \cos \beta - h \sin \alpha + H \cos \alpha + i(G^0 \cos \beta - A \sin \beta) \end{pmatrix}$$

$$\Phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}(G^+ \sin \beta + H^+ \cos \beta) \\ v \sin \beta + h \cos \alpha + H \sin \alpha + i(G^0 \sin \beta + A \cos \beta) \end{pmatrix}$$

- $\tan \beta$ defines basis in Φ space (Higgs basis: $\tan \beta = 0$)
- Higgs-gauge couplings from invariant $s_{\beta-\alpha} \equiv \sin(\beta - \alpha)$
- Parameterisations of potential:
 $\{m_{12}^2, \lambda_{1-7}, \tan \beta\}$ or
 $\{m_{12}^2, m_h, m_H, m_A, m_{H^\pm}, s_{\beta-\alpha}, \lambda_{6-7}, \tan \beta\}$ or ...



Possible additional symmetries

Exact $U(1)_{PQ}$ symmetry

Demanding that the potential has additional $U(1)_{PQ}$ symmetry
 $\Rightarrow m_{12}^2 = 0, \lambda_{5-7} = 0$

(spontaneous breaking gives one more Goldstone boson which after explicit breaking by instanton effects could have given the axion solution to the strong CP-problem)

Exact Z_2 symmetry

Demanding that the potential is symmetric under $\Phi_1 \rightarrow \Phi_1$,
 $\Phi_2 \rightarrow -\Phi_2 \Rightarrow m_{12}^2 = 0, \lambda_{6-7} = 0$

Supersymmetry

Supersymmetry at tree-level \Rightarrow

$$\begin{aligned}\lambda_1 = \lambda_2 &= \frac{g^2 + g'^2}{4}, & \lambda_3 &= \frac{g^2 - g'^2}{4}, & \lambda_4 &= -\frac{g^2}{2}, \\ \lambda_5 = \lambda_6 = \lambda_7 &= 0, & m_{12}^2 &= m_A^2 \cos \beta \sin \beta.\end{aligned}$$



Theoretical constraints

Positivity of potential

Demanding that the potential is bounded from below \Rightarrow

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_3 > -\sqrt{\lambda_1 \lambda_2}$$

If $\lambda_6 = \lambda_7 = 0$: $\lambda_3 + \lambda_4 - |\lambda_5| > -\sqrt{\lambda_1 \lambda_2}$

If $\lambda_6, \lambda_7 \neq 0$: $\lambda_3 + \lambda_4 - \lambda_5 > -\sqrt{\lambda_1 \lambda_2}$
and more complicated constraints

Perturbativity

Cross-section for $2 \rightarrow 2$ Higgs scattering processes $\propto \frac{\lambda_{HHHH}^2}{16\pi^2}$
 \Rightarrow the quartic Higgs couplings λ_{HHHH} cannot be too large for the perturbative series to make sense



Tree-level unitarity

requiring tree-level unitarity for HH and HV_L scattering \Rightarrow limits on eigenvalues of the scattering (S) matrices

$$16\pi S_{(2,1)} = \begin{pmatrix} \lambda_1 & \lambda_5 & \sqrt{2}\lambda_6 \\ \lambda_5 & \lambda_2 & \sqrt{2}\lambda_7 \\ \sqrt{2}\lambda_6 & \sqrt{2}\lambda_7 & \lambda_3 + \lambda_4 \end{pmatrix}$$

$$16\pi S_{(2,0)} = \lambda_3 - \lambda_4$$

$$16\pi S_{(0,1)} = \begin{pmatrix} \lambda_1 & \lambda_4 & \lambda_6 & \lambda_6 \\ \lambda_4 & \lambda_2 & \lambda_7 & \lambda_7 \\ \lambda_6 & \lambda_7 & \lambda_3 & \lambda_5 \\ \lambda_6 & \lambda_7 & \lambda_5 & \lambda_3 \end{pmatrix}$$

$$16\pi S_{(0,0)} = \begin{pmatrix} 3\lambda_1 & 2\lambda_3 + \lambda_4 & 3\lambda_6 & 3\lambda_6 \\ 2\lambda_3 + \lambda_4 & 3\lambda_2 & 3\lambda_7 & 3\lambda_7 \\ 3\lambda_6 & 3\lambda_7 & \lambda_3 + 2\lambda_4 & 3\lambda_5 \\ 3\lambda_6 & 3\lambda_7 & 3\lambda_5 & \lambda_3 + 2\lambda_4 \end{pmatrix}$$



Yukawa sector

General Yukawa couplings for SM fermions with mass eigenstates in flavour vectors D , U , L and ν (neutrinos massless)

$$\begin{aligned}
 -\mathcal{L}_Y = & \overline{D} \frac{\kappa^D s_{\beta-\alpha} + \rho^D c_{\beta-\alpha}}{\sqrt{2}} Dh + \overline{D} \frac{\kappa^D c_{\beta-\alpha} - \rho^D s_{\beta-\alpha}}{\sqrt{2}} DH + i \overline{D} \gamma_5 \frac{\rho^D}{\sqrt{2}} DA \\
 & + \overline{U} \frac{\kappa^U s_{\beta-\alpha} + \rho^U c_{\beta-\alpha}}{\sqrt{2}} Uh + \overline{U} \frac{\kappa^U c_{\beta-\alpha} - \rho^U s_{\beta-\alpha}}{\sqrt{2}} UH - i \overline{U} \gamma_5 \frac{\rho^U}{\sqrt{2}} UA \\
 & + \overline{L} \frac{\kappa^L s_{\beta-\alpha} + \rho^L c_{\beta-\alpha}}{\sqrt{2}} Lh + \overline{L} \frac{\kappa^L c_{\beta-\alpha} - \rho^L s_{\beta-\alpha}}{\sqrt{2}} LH + i \overline{L} \gamma_5 \frac{\rho^L}{\sqrt{2}} LA \\
 & + \left[\overline{U} \{ V_{CKM} \rho^D P_R - \rho^U V_{CKM} P_L \} DH^+ + \overline{\nu} \rho^L P_R LH^+ + \text{h.c.} \right]
 \end{aligned}$$

- κ^F and ρ^F 3×3 matrices: $\kappa^F \equiv \sqrt{2} \frac{M^F}{v}$, ρ^F free (symmetric)
- $P_{R/L} = (1 \pm \gamma_5)/2$
- Non-diagonal $\rho \Rightarrow$ non-MFV CC and FCNC
- Avoided (Glashow & Weinberg) by imposing Z_2 symmetry on Φ_1 , Φ_2 and U_R , D_R , L_R such that each fermion type only couples to one Higgs doublet
 $\Rightarrow \rho^F = \kappa^F \cot \beta$ or $\rho^F = -\kappa^F \tan \beta$
four different types of 2HDM (I, II, III, IV)



Partial decay widths

- $H \rightarrow ff'$ with optional (N)LO QCD corrections
- $H \rightarrow gg$ with optional LO QCD corrections
- $H \rightarrow HH$
- $H \rightarrow HV^*$ including off-shell vector bosons
- $H \rightarrow VV^*$ including off-shell vector bosons
- $H \rightarrow \gamma\gamma$
- $t \rightarrow H^+ b$

here $H = \{h, H, A, H^\pm\}$, $V = \{Z, W\}$


Experimental constraints

- Oblique parameters: contribution to S , T , U , V , W , X compared to SM with Higgs mass m_h^{ref}
- Muon anomalous magnetic moment: 2HDM contribution
- Charged Higgs mass limits from LEP
- Additional Higgs mass limits via HiggsBounds or NMSSMTools (optional)
- Flavour limits from SuperIso (optional)

Example:

2σ limits on Higgs mass differences from S , T , and U with $m_S^2 = m_H^2 s_{\beta-\alpha}^2 + m_h^2 c_{\beta-\alpha}^2$

F. Mahmoudi and O. Stål,
Phys. Rev. D **81** (2010) 035016





Usage/setting parameters/input-output

Programming features


- object oriented code (C++)
- modular structure
- “ready to compile” commandline type programs
- library mode which can be called by user program

Getting started

- download code, manual, and full class documentation from
<http://www.isv.uu.se/thepl/MC/2HDMC>
- system requirements
 - gcc compiler (3.4 and 4 tested)
 - GNU Scientific Library (GSL)
 - HiggsBounds (optional)
 - NMSSMTools (optional)
 - SuperIso (optional)
- adapt makefile and make
- test with Demo-program



Structure of 2HDMC code





LUND
UNIVERSITY

2HDMC

Johan Rathsman

2HDM

TH constraints

Yukawa sector

EX constraints

Usage

Example

rathsman@gungner: ~/private/div_programs/2hdmc\$./CalcPhys 115 250 220 300 0.5 0. 0. 5000 5 2 out_file

2HDM parameters in physical mass basis:

```
m_h:    115.00000
m_H:    250.00000
m_A:    220.00000
m_H+:   300.00000
sin(b-a): 0.50000
lambda_6: 0.00000
lambda_7: 0.00000
m12^2:   5000.00000
tan(beta): 5.00000
```

2HDM parameters in generic basis:

```
lambda_1: 4.15907
lambda_2: 0.68666
lambda_3: 4.63597
lambda_4: -1.74187
lambda_5: -0.36949
lambda_6: 0.00000
lambda_7: 0.00000
m12^2:   5000.00000
tan(beta): 5.00000
```

Tree-level unitarity 1

Perturbativity 1

Stability 1

Mass constraints 1

Charged Higgs 1 (HpHp:1 HpHptau:1 HpHpcS:1)

Oblique parameters:

```
S          1.86057e-02
T          1.56508e-01
U          2.80514e-03
V          4.16485e-03
W          2.39736e-03
X          -6.57052e-05
Delta_rho  1.22358e-03
Delta_amu  3.11201e-11
```

rathsman@gungner: ~/private/div_programs/2hdmc\$



Summary

- General two Higgs Doublet Models – no CP-violation (yet)
- Choice of parameterisations of potential
- Tree-level Higgs masses
- Arbitrary Yukawa sector or “types”
- Yukawa coupling with running quark masses
- Theoretical constraints (positivity, unitarity, perturbativity)
- Electroweak precision tests - oblique parameters, muon $g - 2$
- mass-limits (optionally from HiggsBounds, NMSSMTools)
- Flavour observables from SuperIso
- Partial widths for two-body Higgs decays and non-standard top decays
- Les Houches style input/output
- Madgraph/MadEvent model

D. Eriksson, JR and O. Stål, Comp. Phys. Comm. **181** (2010) 189; 833
<http://www.isv.uu.se/thepl/MC/2HDMC>



LUND
UNIVERSITY

2HDMC

Johan Rathsman

2HDM

TH constraints

Yukawa sector

EX constraints

Usage

Example

backup



Type				
	I	II	III	IV
ρ^D	$\kappa^D \cot \beta$	$-\kappa^D \tan \beta$	$-\kappa^D \tan \beta$	$\kappa^D \cot \beta$
ρ^U	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$
ρ^L	$\kappa^L \cot \beta$	$-\kappa^L \tan \beta$	$\kappa^L \cot \beta$	$-\kappa^L \tan \beta$



LUND
UNIVERSITY

2HDMC

Johan Rathsman

2HDM

TH constraints

Yukawa sector


EX constraints

Usage

Example

Example: Flavour constraints on general 2HDMs



F. Mahmoudi and O. Stål, Phys. Rev. D **81** (2010) 035016





General flavour diagonal couplings

$$\rho^U = \frac{\sqrt{2}}{v} \begin{pmatrix} \lambda_{uu} m_u & 0 & 0 \\ 0 & \lambda_{cc} m_c & 0 \\ 0 & 0 & \lambda_{tt} m_t \end{pmatrix} \quad \text{etc}$$

 $\Delta M_{B_d}:$ $B_u \rightarrow \tau \nu_\tau:$ F. Mahmoudi and O. Stål, Phys. Rev. D **81** (2010) 035016